

Atlasovite

©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Tetragonal (by analogy to nabokoite). *Point Group:* $4/m\ 2/m\ 2/m$. Eight-sided crystals are tabular with prominent {001} modified by {110}, {012} and {014}, to 1 mm; typically intergrown with nabokoite.

Physical Properties: *Cleavage:* On {001}, perfect. *Hardness* = 2–2.5 *D*(meas.) = 4.20(5) *D*(calc.) = 4.12

Optical Properties: Transparent. *Color:* Dark brown; pale gray with yellow internal reflections in reflected light. *Streak:* Pale brown. *Luster:* Vitreous. *Optical Class:* Uniaxial (-). *Pleochroism:* *O* = red-brown; *E* = pale yellow. $\omega = 1.783(3)$ $\epsilon = 1.776(3)$

Cell Data: *Space Group:* $P4/ncc$. $a = 9.86(2)$ $c = 20.58(2)$ $Z = 4$

X-ray Powder Pattern: Tolbachik volcano, Russia. 10.41 (10), 10.75 (9), 2.446 (8), 3.431 (7), 2.890 (7), 4.57 (5), 7.14 (4)

Chemistry:	(1)	(2)
SO ₃	32.21	31.65
TeO ₂	1.29	
V ₂ O ₃	0.68	
Bi ₂ O ₃	12.82	18.42
Fe ₂ O ₃	6.26	6.31
CuO	38.15	37.73
ZnO	1.02	
PbO	2.35	
K ₂ O	3.85	3.72
Cs ₂ O	0.01	
Cl	2.92	2.80
-O = Cl ₂	0.66	0.63
Total	100.90	100.00

(1) Tolbachik volcano, Russia; by electron microprobe, average of four analyses, total Fe as Fe₂O₃; corresponding to K_{1.01}(Cu_{5.90}Zn_{0.15})_{Σ=6.05}(Fe_{0.97}V_{0.11})_{Σ=1.08}(Bi_{0.68}Pb_{0.12}Te_{0.10})_{Σ=0.90}O_{4.19}(SO₄)_{4.95}Cl_{1.01}. (2) KCu₆FeBiO₄(SO₄)₅Cl.

Polymorphism & Series: Forms a series with nabokoite.

Occurrence: As a volcanic sublimate.

Association: Nabokoite, anglesite, dolerophanite, euchlorine, chloroxiphite, atacamite, piypite, chalcocyanite, alarsite, fedotovite, lammerite, klyuchevskite, langbeinite, hematite, tenorite.

Distribution: From the Tolbachik fissure volcano, Kamchatka Peninsula, Russia.

Name: For Vladimir Vasil'evich Atlasov (1661/1664–1711), Russian traveller who first explored the Kamchatka Peninsula.

Type Material: A.E. Fersman Mineralogical Museum, Academy of Sciences, Moscow, Russia.

References: (1) Popova, V.I., V.A. Popov, N.S. Rudashevskiy, S.F. Glavatskikh, V.O. Polyakov, and A.F. Bushmakina (1987) Nabokoite Cu₇TeO₄(SO₄)₅•KCl and atlasovite Cu₆Fe³⁺Bi³⁺O₄(SO₄)₅•KCl. New minerals of volcanic exhalations. *Zap. Vses. Mineral. Obshch.*, 116, 358–367 (in Russian with English abs.). (2) (1988) *Amer. Mineral.*, 73, 927–928 (abs. ref. 1). (3) Pekov, I.V. (1998) Minerals first discovered on the territory of the former Soviet Union. *Ocean Pictures*, 30.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.