


Harkerite

©2001 Mineral Data Publishing, version 1.2

Crystal Data: Hexagonal, pseudocubic. *Point Group:* $\bar{3} 2/m$. As simple pseudo-octahedral crystals, to 5 mm. *Twinning:* Observed, with four rhombohedral individuals each ideally oriented || to one of the pseudocubic 3-fold axes.

Physical Properties: Hardness = n.d. D(meas.) = 2.95–2.96 D(calc.) = [3.02]

Optical Properties: Semitransparent. *Color:* Colorless. *Luster:* Vitreous.

Optical Class: Uniaxial, but typically sensibly isotropic; sections may separate into four fields with weak birefringence in the diagonal position. $n = 1.649$ – 1.653

Cell Data: *Space Group:* $R\bar{3}m$. $a = 10.439$ $c = 51.303$ $Z = 3$

X-ray Powder Pattern: Broadford area, Isle of Skye, Scotland.

2.61 (vs), 5.22 (s), 2.13 (s), 1.84 (s), 1.51 (s), 3.39 (m), 3.01 (m)

Chemistry:	(1)	(2)	(3)
SiO ₂	14.17	16.82	16.37
B ₂ O ₃	7.77	7.0	7.11
Al ₂ O ₃	2.84	3.13	3.47
Ce ₂ O ₃		0.8	
Fe ₂ O ₃	0.85	1.78	
FeO	0.46		
MnO	0.02	0.13	
MgO	11.15	10.01	10.98
CaO	46.23	44.13	45.85
SrO		0.2	
F		0.15	
Cl	1.36	0.06	
H ₂ O ⁺	0.81	1.25	1.23
H ₂ O ⁻	0.11		
CO ₂	14.94	15.07	14.99
–O = (F, Cl) ₂	0.31	0.07	
Total	100.40	100.46	100.00

(1) Camas Malag, Isle of Skye, Scotland. (2) Alban Hills, Italy; by electron microprobe, spectrophotometry, elemental analyzer, and mass spectrometer. (3) Ca₂₄Mg₈Al₂(SiO₄)₈(BO₃)₆(CO₃)₁₀•2H₂O.

Occurrence: In skarns in dolomitic limestones at the contact with granite (Camas Malag, Scotland); in volcanic ejecta (Alban Hills, Italy).

Association: Calcite, monticellite, diopside, magnetite, vesuvianite, cuspidine, phlogopite.

Distribution: At Camas Malag, near Broadford, Isle of Skye, Scotland. On the Chersky Ridge, Tas-haiatah, northeastern Siberia, Russia. From the Alban Hills, Lazio, Italy. Large crystals at Nordmark, Sweden.

Name: For Alfred Harker (1859–1939), British petrologist, Cambridge University, Cambridge, England.

Type Material: n.d.

References: (1) Tilley, C.E. (1951) The zoned contact-skarns of the Broadford area, Skye: a study of boron-fluorine metasomatism in dolomites. *Mineral. Mag.*, 29, 621–667. (2) (1952) *Amer. Mineral.*, 37, 359 (abs. ref. 1). (3) Ostrovskaya, I.V., N.N. Pertsev, and I.B. Nikitina (1966) Sakhaite, a new carbonate-borate of calcium and magnesium. *Zap. Vses. Mineral. Obsch.*, 95, 193–202 (in Russian). (4) Giuseppetti, G., F. Mazzi, and C. Tadini (1977) The crystal structure of harkerite. *Amer. Mineral.*, 62, 263–272. (5) Barbieri, M., C. Cozzupoli, M. Federico, M. Fornaseri, S. Merlino, P. Orlandi, and L. Tolomeo (1977) Harkerite from the Alban Hills, Italy. *Lithos*, 10, 133–141.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.